

SUCKING CHEST WOUND

30 Second Review RN, LPN

Sucking Chest Wound

DEF: A chest wall defect that allows air to enter the pleural space.

S/S: Acute respiratory distress, diminished or absent breath sounds.

RX: Arrange transport. While waiting apply petrolatum dressing on 3 sides, start high flow Oxygen 15L mask, and may start 1-2 large bore IV's TKO, and obtain EKG.

SKILL LEVEL: RN, LPN

DEFINITION: A chest wall defect that allows air to enter the pleural space causing respiratory distress.

DATA BASE:

Subjective: Describe mechanism of injury i.e., penetrating chest trauma. Elicit time, place, source, and description of pain.

Objective:

- Note a sucking noise as air enters and leaves the chest through the wound.
- Acute respiratory distress.
- Diminished or absent breath sounds on the afflicted side.
- Possible subcutaneous emphysema.

Assessment: Sucking chest wound.

Plan: **These patients all require transport to an emergency facility. Have somebody arrange transportation while the following are being done.**

1. Apply petrolatum dressing or plastic wrap dressing then cover with bulky dressing. ***The dressing should be taped on three sides. A completely occlusive dressing could make the situation worse.***
2. Maintain patient's airway, breathing, and circulation.
3. Administer Oxygen, high flow 15L by mask.

Sucking Chest Wound

4. Do not remove any object protruding from the chest as this will destroy pressure gradient even faster and increase bleeding.
5. Establish IV therapy with large bore catheter, two sites if possible.
6. May notify physician while awaiting transport.

APPROVED:

Health Services Manager

Date

Chief Medical Officer

Date



Medical Director

3/2/09

Date

Effective Date: _____

Revised: January 2009